



Therapeutic Potential of Orally Administered Aqueous Seed Extract of *Garcinia kola* on Experimental Colitis Induced by Acetic Acid Colitis in Wistar Rats

A. Perede ^{a*}, O. G. Aviowro ^b, M. S. Abubakar ^c,
S. M. Sahabi ^d, I. Igbinosa ^a, S. M. Gamde ^a, U. Abubakar ^a,
M. N. Chukwu ^a and M. Bello ^e

^a Department of Histopathology, School of Medical Laboratory Science, Usmanu Danfodiyo University, P.M.B. 2346, Sokoto, Nigeria.

^b Department of Medical Laboratory Science, Delta State University of Nigeria, Nigeria.

^c Department of Pathology Faculty of Veterinary Medicine, Usmanu Danfodiyo University, P.M.B. 2346, Sokoto, Nigeria.

^d Department of Morbid Anatomy and Forensic Medicine, Usmanu Danfodiyo University, P.M.B. 2346, Sokoto, Nigeria.

^e Department of Chemical Pathology, Medical Laboratory Science, Specialist Hospital Sokoto, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajrb/2024/v14i6322>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116402>

Original Research Article

Received: 20/03/2024

Accepted: 24/05/2024

Published: 03/10/2024

*Corresponding author: E-mail: histospirit@gmail.com;

ABSTRACT

Introduction: Erosive colitis is a common disorder of the gastrointestinal tract with unknown cause, characterized by recurrent erosion of the colonic mucosal lining.

The pathogenesis likely involves genetic, environmental, and immunologic factors.

Aim: This study investigated the therapeutic effects of *Garcinia kola* (GK) on acetic acid-induced erosive colitis in adult male Wistar rats. Erosive colitis (EC) was induced in male Wistar rats by intra-rectal administration of 2 ml of 4 % acetic acid at 8 cm proximal to the anus for 30 seconds.

Methodology: Thirty-six animals were divided into six groups (n=6), negative control (no erosion, no GK); positive control (erosive colitis without treatment of GK extract); treated with 100, 200, 400mg/kg GK, standard treatment group (Prednisolone) for 14 days.

Results: *Garcinia kola* significantly improved Erosive colitis-induced reduction in mean stool consistency and mean macroscopic erosion score ($P < 0.05$). Injection of GK also significantly reduced the mean microscopic erosion score when compared to untreated EC control ($P < 0.01$). GK reduced the colonic mucosa injury of the Wistar rat.

Conclusion: Results of this study provide scientific evidence that lends credence to the use of aqueous seed extract of *Garcinia kola* in folklore medicine in the management of colitis erosion.

Keywords: Erosive colitis; *Garcinia Kola*; wistar rats.

1. INTRODUCTION

“Erosive colitis is a disease condition in which the colon loses its structural integrity and the ability to absorb water and form faeces” [1]. “Patients with erosive colitis may present with symptoms such as watery diarrhoea, abdominal pain, tenesmus, urgency, fever, subjective fatigue, or blood in the stool” [2]. “There are several different causes of erosive colitis, including infection, autoimmunity, and ischemia due to blood supply, toxin exposure, immunodeficiency, and radiation exposure” [3]. “Erosive colitis is a common and increasing disease incidence worldwide. Nearly one million individuals in the United States and Europe are affected by this condition and many more globally. Over the past decade, since the publication of the last guideline from the American College of Gastroenterology (ACG) on this topic, it occurs throughout the world but is more common in urban areas and present in teens in early 20s” [4]. “Despite the fact that cause of erosive colitis remains poorly understood, complex interactions among genetic, environmental, immunological, and reactive oxygen species (ROS) have been implicated in the pathogenesis of erosive colitis” [5,6].

“*Garcinia kola* is a species of flowering plant that belongs to a family of tropical plants known as Clusiaceae or Guttiferae also known as the African wonder nut. In Nigerian languages, it is commonly called *Namijin goro* in Hausa, *Agbilu* in Igbo, and *Orogbo* in Yoruba. *Garcinia kola* has economic and cultural values across West and

Central African countries where the nuts are commonly chewed and used for traditional ceremonies” [7]. “The seeds are also used in folk medicine in many herbal formulations and have potential therapeutic benefits due largely to the activity of their flavonoids and other bioactive compounds” [8]. “This plant has been referred to as a “wonder plant” because every part of it has been found to be of medicinal importance. The seeds are chewed as an aphrodisiac or used to cure cough, dysentery, chest colds, liver disorders, diarrhoea, laryngitis, bronchitis, and gonorrhea” [9]. “The seed is used to prevent and relieve colic; it can also be used to treat headaches, stomach aches and gastritis” [10].

1.1 Statement of the Problem

“Erosive colitis is a common gastrointestinal disorder. Although there is little epidemiologic data conducted in developing countries like Nigeria it has an increasing incidence worldwide. Nearly one million individuals in the United States and Europe are affected by this condition and many more globally. Over the past decade, since the publication of the last guideline from the American College of Gastroenterology (ACG) on this topic, epidemiological studies from all over the world have stated that the incidence and prevalence of erosive colitis are increasing with time and in different regions around the world indicating its emergence as a global disease” [11].

1.2 Justification of the Study

“Acetic acid-induced erosive colitis is a commonly employed and easily inducible model of erosion. Intra rectal administration of a dilute solution of acetic acid causes non-transmural erosion characterized by increased neutrophil infiltration into the intestinal tissue, massive necrosis of mucosal and sub-mucosal layers, vascular dilation, oedema and sub-mucosal erosion that are noteworthy features of human erosive colitis” [12]. The characteristic feature is an imbalance between oxidant and antioxidant substances [13].

“Acetic acid-induced erosive colitis in the colon of Wistar rats bears close resemblance to human erosive colitis in terms of pathogenesis, histopathological features, and inflammatory mediator profile and is therefore a reliable animal model that can be useful for the evaluation of drugs for erosive colitis” [14].

“Extract of *Garcinia kola* has been reported to have a myorelaxant effect in addition to anti-inflammatory action on animal models of erosive colitis” [15].

2. METHODS

2.1 Study Location

The study was conducted at the Histopathology Department of the School of Medical Laboratory Sciences, Usmanu Danfodiyo University, Sokoto, Sokoto State

2.2 Experimental Animals

Adult Male Wistar rats (155-204g) were procured from the animal house of the Faculty of Pharmaceutical Sciences, Usmanu Danfodiyo University, Sokoto. They were kept in a well-ventilated room with optimum environmental conditions of temperature, relative humidity, and dark/light cycle and were fed standard feed pellets and tap water ad libitum. They were acclimatized for two weeks prior to the experiment.

2.3 Plant Collection

Seeds of *Garcinia kola* were purchased from Sokoto Central Market (Shehu Shagari Gate) Sokoto, Nigeria. The plant material was subjected to identification and authentication at the Department of Botany, Usmanu Danfodiyo University, where a voucher number was

allocated a voucher number –PCG/ UDUS/ GUH/0001 and deposited at the herbarium.

2.4 Extract Preparation

“The seed was cleaned and air-dried at room temperature for 7 days and ground to a fine powder using mortar and pestle. Five hundred (500) grams of the powdered material was macerated in 1.5 L of distilled water and left for 24 hours after which it was filtered using Whitman's filter paper. The filtrate was dried in a hot air oven at 40°C to give 39.8g of the aqueous seed extract which was used for the study. The percentage yield was calculated to be 7.96% and the dried extract was stored in an airtight container” [16].

2.5 Acute Toxicity Testing

Acute toxicity testing was conducted using [17]. In Phase I, nine rats were used and randomly assigned into 3 groups of 3 rats each. The 1st group was administered 10mg/kg body weight of the extract using an oral cannula, and the 2nd and 3rd groups received 100mg/kg and 1000mg/kg body weight, respectively. The animals were then observed for 24 hours to monitor their behavior for signs of toxicity as well as mortality. In Phase II, three rats were used and randomly placed into 3 groups of an animal each. The animals were administered high doses of 1600mg/kg, 2900mg/kg, and 5000mg/kg, respectively. They were then observed for 24 hours for signs of toxicity and mortality.

2.6 Colitis Induction

“All animals (except group I) fasted for 6 hours prior to the study, water ad libitum, and given mild anesthesia before induction of erosive colitis and 2ml acetic acid (4% v/v) in 0.9% saline were infused for 30s using a soft flexible pediatric catheter size of 6F 2mm in diameter, inserted through the rectum into the colon up to a distance of 8cm and maintained in a supine Trendelenburg position for 30 seconds to prevent leakage of the intracolonic instillation” [18].

2.7 Experimental Design

The study is an experimental which consists of Thirty six (36) wistar rats randomly divided in to six (6) groups as shown in the table below:

Table 1. Summary of experimental design

Experimental group	Induction of EC	Treatment given	Duration of treatment
1 (negative control) (5 rats)	Distilled water	No treatment given	14 days
2 (positive control) (5 rats)	2mls(4% acetic acid)	No treatment given	14 days
3 (treatment group 1) (5 rats)	2mls(4% acetic acid)	100mg/kg(A.S.E.G.K)	14 days
4 (treatment group 2) (5 rats)	2mls(4% acetic acid)	200mg/kg(A.S.E.G.K)	14 days
5 (treatment group 3) (5 rats)	2mls(4% acetic acid)	400mg/kg(A.S.E.G.K)	14 days
6 (prednisolone group 4) (5 rats)	2mls(4% acetic acid)	2mg/kg	14 days

Table 2. Scoring based on stool consistency

Score	Stool Consistency
0	Normal stool
1	Soft, stool but still formed
2	Soft, wet stool but unformed
3	Soft, wet stool + blood
4	Bloody diarrhea

Table 3. Scoring based on macroscopic characteristics

Score	Macroscopic changes
0	No macroscopic change
1	Mucosal erythema, and hyperemia at the sites
2	Mild mucosal oedema, slight bleeding, or small erosions
3	Moderate mucosal oedema, slight bleeding erosions
4	Severe erosion, oedema, and tissue necrosis

Table 4. Scoring of ulcer area

Score	Changes
0	Normal coloured colon
0.5	Red colouration
1	Spot erosion
1.5	Haemorrhagic streaks
2	Erosion ≥ 3 but ≤ 5
3	Erosion >5

Table 5. Histological scoring pattern

Score	Histological Changes
0	No abnormality detected
1 (mild)	Damage / active changes up to 25%
2 (moderate)	Damage / active changes more than 25% but less than 50 %
3 (severe)	Damage / active changes more than 50 %

2.8 Scoring and Assessment

Scoring based on stool consistency: Every morning, stool from all rats from the six groups was examined physically and then Scored. The following scoring pattern of [19] was used.

Scoring based on macroscopic characteristics: Pieces of rat colon (10 cm long each) were scored for macroscopic features using the scoring pattern as shown in

the Table 3 Above as described by Deshmukh [19].

Scoring of ulcer area: The erosion area was determined by the method described by Dengiz [20].

Using the method of [21], "the histomorphological parameters analyzed were erosive cell infiltrate (for severity and extent), epithelial changes (to assess hyperplasia and erosion), and mucosal

architecture (for goblet cell loss and altered crypts)".

2.9 Data and Statistical Analysis

All the results were expressed as mean $\pm$ S.D. Data analysis was performed using

GraphPad Prism 6.0 software (GraphPad, San Diego, USA). Statistical comparison between drug-treated groups and colitis control animals was done using one-way ANOVA. A value of $p < 0.05$ was considered to be statistically significant.

3. RESULTS

Table 6. The physical properties of aqueous seed extract of *Garcinia kola*

Plant Part	Extract Type	% Yield	Texture	Colour	Smell
Seed	Aqueous	7.96%	Soft gel	Brown	Sweet

Table 7. Acute toxicity of aqueous seed extract of *Garcinia kola* Wistar Rats (N=12)

Dose	Mortality	
	Phase I	Phase II
10mg	0/3	-
100mg	0/3	-
1000mg	0/3	-
1600mg	-	0/1
2900mg	-	0/1
5000mg	-	0/1




Plate 1A. Photomicrograph of Colonic Tissue from Control Animal. Intact colonic mucosa from control Animals showing normal crypts (black arrow), goblet cells (red arrow), submucosa (white arrow), and muscularis propria (green arrow) (H&E. Mag x 100)




Plate 1B. Photomicrograph of Colonic Tissue from Colitis Control Group. The section shows oedema (black arrow), numerous inflammatory cells extending to the submucosa (black arrow), distortion of crypts and goblet cells and mucosal erosion (red arrow), and mucosal erosion (white arrow) (H&E. Mag x 100)




Plate 1C. Photomicrograph of Colonic Tissue from Animal Receiving 100mg/kg of the Extract. The section shows mild crypt distortion (black arrow) with a few goblet cells (red arrow). Inflammatory cells are seen in the mucosa and submucosa (white arrow) (H&E. Mag x 100)




Plate 1D. Photomicrograph of Animals receiving 200mg/kg of the Extract Section shows almost normal crypts (black arrow) with normal goblet cell numbers (red arrow). Infiltrates cells are seen mostly within the mucosa and submucosa (white arrow) (H&E. Mag x 100)




Plate 1E. Photomicrograph of Animal Receiving 400mg/kg of the Extract. The section shows normal goblet cells (black arrow) and normal crypts (red arrow). A clearing of inflammatory cells and polymorphs are seen in the mucosa (white arrow). (H&E. Mag x 100)

Table 8. Effect of A. S. E. G. K. on some physical parameters

Parameters	Normal	Acetic acid		Garcinia kola			P-Values
		Control	100mg/kg	200mg/kg	400mg/kg	2mg/kg	
SC	1.00 ± 0.00	3.00 ± 0.00	1.50 ± 0.29	1.25 ± 0.25	1.25 ± 0.25	1.25 ± 0.25	0.000
MIC	0.00 ± 0.00	3.75 ± 0.25	2.75 ± 0.25	2.25 ± 0.25	1.25 ± 0.25	0.25 ± 0.25	0.000
MAC	0.00 ± 0.00	3.75 ± 0.25	2.75 ± 0.25	2.25 ± 0.25	1.25 ± 0.25	0.25 ± 0.25	0.000

Keynote: Data are expressed as Mean±SD. SC- Stool consistency, MIC- Microscopic characteristic, and MAC- Macroscopic characteristics

Table 9. Effect of A. S. E. G. K. on macroscopic colonic features

Parameters	Normal	Acetic acid		G. kola			P-Values
		Control	100mg/kg	200mg/kg	400mg/kg	2mg/kg	
MAC	0.00 ± 0.00	3.75 ± 0.25	2.75 ± 0.25	2.25 ± 0.25	1.25 ± 0.25	0.25 ± 0.25	0.000

Keynote: Data are expressed as Mean±SD

Table 10. Effect of A.S.E.G.K. on colon histological parameters

Parameter	normal	Acetic acid		G. kola		Prednisolone	P-Values
		Control	100mg/kg	200mg/kg	400mg/kg	2mg/kg	
Infiltrates	0.00 ± 0.00	3.75 ± 0.259	3.25 ± 0.25	2.25 ± 0.25	1.75 ± 0.48	0.75 ± 0.25	0.000
Severity							
Epithelial	0.25 ± 0.10	3.75 ± 0.25	3.00 ± 0.00	2.25 ± 0.29	1.50 ± 0.29	0.50 ± 0.29	0.000
Hyperplasia							
Goblet cell loss	0.00 ± 0.00	3.75 ± 0.25	3.25 ± 0.25	2.50 ± 0.25	1.50 ± 0.29	0.75 ± 0.25	0.000
Crypts	0.00 ± 0.00	3.75 ± 0.25	3.25 ± 0.25	2.25 ± 0.25	1.50 ± 0.29	0.75 ± 0.25	0.000
alteration							

Keynote: Data are expressed as Mean±SD




Plate 2. Photomicrograph of Colonic Tissue from Treatment Control Group. The section shows a normal crypt (black arrow) with an adequate amount of goblet cells (red arrow). Polymorphs are scanty (white arrow). There is a clearing of most infiltrates by Prednisolone. (H&E. Mag x 100)

4. DISCUSSION

“The extraction yield is a measure of the solvent efficiency to extract specific components from an original material. The aqueous extraction method used in this research produced a percentage yield of 7.96%. This is in agreement with the work” [22]. The extract was brown, with a crystalline shine and a slightly sweet smell [23]. Research by Somboro [22] has shown that several compounds are absent or present in different quantities from plants from different provenances, indicating the presence of chemotypes. This may account for the similar results in the percentage yield of the crude extract.

In the acute toxicity study, all the graded doses up to 5000mg/kg showed no sign of toxicity in the animal and no mortality was recorded in the study. The LD50 of aqueous seed extract of *Garcinia kola* was found to be higher than 5000mg/kg body weight. This result is consistent with the findings of [24], [25,26,27,28,29].

“Treatment of erosive colitis rats with the aqueous seed extract of *Garcinia kola* in this study reduced erosion scores. *Garcinia kola* contains biflavonoids and bioflavonoids have

been shown to stimulate angiogenesis, an important facet of tissue healing” [30].

“It was observed that animals receiving the aqueous seed extract of *Garcinia kola* showed improvement in stool consistency in a dose-dependent manner. It was also seen in animals receiving the extract but improved remarkably in a dose-dependent manner. Animals receiving the highest dose of the extract showed comparatively similar results to normal control groups. It was also observed that *Garcinia kola* extract ameliorated diarrhoea score” [31]. “Swelling is one of the symptoms of inflammation. The result of the study indicated that treatment of erosive colitis rats with aqueous seed extract of *Garcinia kola* decreased tissue thickness. Histopathological damage was evaluated in colonic samples stained with haematoxylin and eosin. In normal animals, epithelial crypts of the mucosal layer were intact. There was no infiltration of inflammatory cells. The intra-rectal instillation of acetic acid resulted in and significant development of transmural necrosis, submucosal oedema, erosion along with cellular infiltration, and loss of epithelial crypts and goblet cells” [24]. Animals fed with the extract showed a reduction in the extent of damage in a dose dependent manner) 100mg/kg, 200mg/kg, and animals receiving the

control drug - Prednisolone (2mg/kg). Animals fed with the highest dose 400mg/kg showed clearing of inflammatory cells with decreased goblet cells. This result is consistent with the findings of Fiot et al. [24], and [25,26,27,28,29].

The histological studies of erosive colitis-treated rats reveal regenerating mucosal layer with regenerating colonic crypts and goblet cells (Plate 1A and Plate 1B) in contrast to the histology of colitis rats which are characterized by indistinguishable colonic layers (Plate 2). This result is consistent with the findings of [24], and [25,26,27,28,29].

“Microscopically, mucosal and sub-mucosal inflammatory cellular infiltrations were detected. It was noted that the animals that received acetic acid showed a significant increase in mucosal and submucosal inflammatory cellular infiltrates compared to the control animals. It also indicated that there was a mild decrease in the severity but a marked improvement in the extent of mucosal and submucosal inflammatory cellular infiltration in animals receiving the extract. 400mg/kg treatment group cleared all polymorphs in the colon when compared to Prednisolone which is a standard drug of choice. This is indicative of the superiority and relevance of the action of a high dose of *Garcinia kola* on the inflammatory cells. More so, Prednisolone which is a standard drug of choice effectively reduced all histological changes such as oedema, crypt distortion, and goblet cell loss and tissue injury by virtue of its healing properties” [16,32].

5. CONCLUSIONS

The present study showed that treatment of colitis rats with *Garcinia kola* extract decreased erosion score by stimulating colonic healing and reduced diarrhea score and tissue thickness. The extract also improved the weight of erosive colitis rats in 14 days after treatment. Therefore, aqueous seed extract of *Garcinia kola* exhibited a therapeutic effect. Oral administration of aqueous extract of the seed of *Garcinia kola* at doses used in this study showed notable improvements in body weight compared to erosive colitis control animals.

However, the effect of Prednisolone which is a standard drug of choice was not as potent as to 400mg/kg treatment group used in the study. Prednisolone which is a standard drug of choice also showed remarkable improvement in the

scores of both macroscopic and microscopic colonic parameters compared to control groups.

In conclusion, the results of this study provide scientific evidence that lends credence to the use of *Garcinia kola* extract in folklore medicine in the management of colitis erosion.

ETHICAL APPROVAL

Ethical approval for this research was sought from the Ethics Committee on Research and Experiments of Usmanu Danfodiyo University, Sokoto.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Rubin E, Fred G, Roland SD. Rubin Pathology. 4th ed. Philadelphia: 2005;355-366.
2. Molodecky NA, Soon IS, Rabi DM, Ghali WA, Ferris M, Chernoff G, Benchimol EI, Panaccione R, Ghosh S, Barkema HW, Kaplan GG. Increasing incidence and prevalence of the inflammatory bowel diseases with time, based on systematic review. *Gastroenterology*. 2012;142(1):46–54, e42.
3. Yadav S, Dave M, Edakkanambeth Varayil J, Harmsen WS, Tremaine WJ, Zinsmeister AR, Sweetser SR, Melton LJ, Sandborn WJ, Loftus EVA population-based study of incidence, risk factors, clinical spectrum, and outcomes of ischemic colitis. *Clin Gastroenterol Hepatol*. Apr. 2015;13(4):731-8.e1-6;quiz e41
4. Kumar V, Abbas AK, Aster JC. Robbins and cotran's pathologic basis of disease. 9th Edition. Elsevier-Saunders E-book. 2015;Chapter 17 –Pp431-433.
5. Fiocchi C. Inflammatory bowel disease: Etiology and pathogenesis. *Gastroenterology*. 1998;115:182-205.
6. Papadakis KA, Targan SR. Current theories on the cause of inflammatory bowel disease. *Gastroenterology Clinics of North America*. 1999;28:283-96.
7. Eleyinmi AF, Bressler DC, Amoo IA, Sporns P, Oshodi AA. Chemical composition of bitter cola (*Garcinia kola*)

- seed and hulls. Polish Journal of Food and Nutrition Sciences. 2006;15(56):395-400.
8. Akintonwa A, Essien AR. Protective effects of *Garcinia kola* seed extract against paracetamol-induced hepatotoxicity in rats. Journal of Ethnopharmacology. 1990; 29:207–219.
9. Adesina SK, Gbile ZO, Odukoya OA, Akinwusi DD, Illoh HC, Jayeola AA. Survey of indigenous useful plants of West Africa with special emphasis on medicinal plants and issues associated with their management. The United Nations University Programme on natural resources in Africa, 2nd edition. 1995;84-90.
10. Ayensu ES. Medicinal plants of West Africa, Reference Publication Inc; Algonac, Michigan. 1978;162.
11. Bernstein CN, Fried M, Krabshuis JH, Cohen H, Eliakim R, Fedail S, Gearry R, Goh KL, Hamid, S, Khan AG, LeMair AW. World gastroenterology organization practice guidelines for the diagnosis and management of IBD in 2015. Inflammatory Bowel Diseases. 2015;16(1):112-124.
12. Hartmann RM, Morgan-Martins MI, Tieppo J, Fillmann HS, Marroni NP. Effect of *Boswellia serrata* on antioxidant status in an experimental model of colitis rats induced by acetic acid. Digestive Diseases and Sciences. 2012;57:2038-2044.
13. Bitiren M, Karakilcik AZ, Zerir M, Ozardali I, Selek S, Nazligül Y, Ozgonul A, Musa D, Uzunkoy, A. Protective effects of selenium and vitamin E combination on experimental colitis in blood plasma and colon of rats. Biological Trace Element Research. 2010;136:87-95.
14. Randhawa PK, Singh K, Singh N, Jaggi AS. A review on chemical-induced inflammatory bowel disease models in rodents. Korean Journal of Physiology and Pharmacology. 2014;18:279-288.
15. Adaramoye OA, Farombi EO, Adeyemi EO, Emerole GO. Inhibition of human low density lipoprotein oxidation by flavonoids of *Garcinia kola* seeds. Pakistan Journal of Medicine and Science. 2005b;21(3):331-339.
16. Akuodor GC, Essien AD, David-Oku E, Chilaka KC, Akpan JL, Ezeokpo B, Ezeonwumelu JO. C. Gastroprotective effect of the aqueous leaf extract of *Guiera senegalensis* in Albino rats. Asian Pacific Journal of Tropical Medicine. 2013; 6(10):771-775.
17. Lorke D. A new approach to practical acute toxicity testing. Archives of Toxicology. 1983;54:275-287.
18. Fabia R, Willén R, Ar'Rajab A, Andersson R, Ahrén B, Bengmark S. Acetic acid-induced colitis in the rat: A reproducible experimental model for acute ulcerative colitis. European Surgical Research. 1992;24:211-225.
19. Deshmukh CD, Veeresh B, Pawar AT. Protective effect of *Embllica officinalis* extract on acetic acid induced colitis in rats. Journal of Herbal Medicine and Toxicology. 2010;4(2):25-29.
20. Dengiz GO, Gursan N. Effects of *Momordica charantia* L. (Cucurbitaceae) on indomethacin-induced ulcer model in rats. Turkish Journal of Gastroenterology. 2005;16(2):85-88.
21. Erben U, Loddenkemper C, Doerfel K, Spieckermann S, Haller D, Heimesaat MM, Zeitz M, Siegmund B, Kühl AA. A guide to histomorphological evaluation of intestinal inflammation in mouse models. International Journal of Clinical and Experimental Pathology. 2011;7(8):4557-4576.
22. Somboro AA, Patel K, Sidibe L, Chalchat JC, Figueredo G, Ducki S, Troin Y, Chalard P. An ethnobotanical and phytochemical study of the African medicinal plant *Guiera senegalensis* J.F. Gmel. Journal of Medicinal Plants Research. 2014;5(9):1639–1651.
23. Saidu TB, Galadima M, Jigam AA, Abalaka EM. Antibacterial and potential toxicological evaluation of leaf extracts of *Guiera senegalensis* against gram negative bacteria. Journal of Advances in Medical and Pharmaceutical Sciences. 2016;9(1):1-13.
24. Fiot J, Sanon S, Azas N, Mahiou V, Jansen O, Angenot L, Balansard G, Oliver E. Phytochemical and pharmacological study of roots and leaves of *Guiera senegalensis* JF Gmel (Combretaceae). Journal of Ethnopharmacology. 2006;106(2):173 – 178.
25. Sule MS, Mohammed SY. Potency of partially purified anthocyanin from leaf extract of *Guiera senegalensis* against carbon tetrachloride – induced lipoperoxidation in rats. Bayero Journal of Pure and Applied Science. 2009;2(2):155-158.
26. Jigam AA, Akanya HO, Dauda BE, Ogbadoyi EO. Antiplasmodial, analgesic

- and anti-inflammatory effects of crude *Guiera senegalensis* Gmel (Combretaceae) leaf extracts in mice infected with *Plasmodium berghei*. Journal of Pharmacognosy and Phytotherapy. 2011;3(10): 150-154.
27. Shettima YA, Tijjani MA, Karumi Y, Sodipo OA. Phytochemical and anti-diarrhoeal properties of methanolic extracts of *Guiera senegalensis*. Gmel. International Research Journal of Pharmacy. 2012;3(11):61-65.
 28. Mohammed SY, Sule MS. Potency of partially purified anthocyanin from leaf extract of *Guiera senegalensis* against carbon tetrachloride – induced lipoperoxidation in rats. Bayero Journal of Pure and Applied Sciences. 2009;2(2): 155–158.
 29. Bako HY, Ibrahim M, Mohammad JS, Zubairu M, Bulus T. Toxicity studies of aqueous, methanolic and hexane leaf extracts of *Guiera senegalensis* in rats. International Journal of Scientific & Engineering Research. 2014;5(10):1338-1347.
 30. Kilicoglu SS, Bulent K, Ezra E. Ultrastructural view of colon anastomosis under Propolis effect by transmission electron microscopy. World Journal of Gastroenterology. 2008;14(30):4763-4770.
 31. Thiagarajah JR, Verkman AS. CFTR inhibitors for treating diarrheal disease. Clinical Pharmacology and Therapeutics. 2012;92(3):287-90.
 32. Al Shafei NK, Ahmed SM, Nour AY. Preliminary observations on the uses of *Guiera senegalensis* as a traditional medicinal plants in Western Kordufan, Sudan. International Journal of Applied and Pure Science and Agriculture. 2016a; 2(5):42–48.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/116402>